

AD-753 610

THE INVERSE OF A TRIDIAGONAL MATRIX

Palmer R. Schlegel, et al

Ballistic Research Laboratories
Aberdeen Proving Ground, Maryland

September 1972

DISTRIBUTED BY:

NTIS

National Technical Information Service
U. S. DEPARTMENT OF COMMERCE
5285 Port Royal Road, Springfield Va. 22151

BRLR 1612

AD 753610

BRL


AD

REPORT NO. 1612

THE INVERSE OF A TRIDIAGONAL MATRIX

by

Palmer R. Schlegel
William Clare Taylor



September 1972

Approved for public release; distribution unlimited.

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
U.S. Department of Commerce
Springfield, VA 22151

BALLISTIC RESEARCH LABORATORIES
ABERDEEN PROVING GROUND, MARYLAND

Destroy this report when it is no longer needed.
Do not return it to the originator.

Secondary distribution of this report by originating or
sponsoring activity is prohibited.

Additional copies of this report may be purchased from
the U.S. Department of Commerce, National Technical
Information Service, Springfield, Virginia 22151

ACQUISITION BY	
HTIC	WHIS Section <input checked="" type="checkbox"/>
DOC	Self Section <input type="checkbox"/>
UNCLASSIFIED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY	
DATE	
A	

The findings in this report are not to be construed as
an official Department of the Army position, unless
so designated by other authorized documents.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION	
Ballistic Research Laboratories Aberdeen Proving Ground, Maryland 21005		Unclassified	
3. REPORT TITLE		2b. GROUP	
THE INVERSE OF A TRIDIAGONAL MATRIX		N/A	
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
BRL Report			
5. AUTHOR(S) (First name, middle initial, last name)			
Palmer R. Schlegel, William Clare Taylor			
6. REPORT DATE		7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
September 1972		1711	1
8a. CONTRACT OR GRANT NO.		8a. ORIGINATOR'S REPORT NUMBER(S)	
b. PROJECT NO. 1T062110A027		Report No. 1612	
c.		8b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.			
10. DISTRIBUTION STATEMENT			
Approved for public release; distribution unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
		U.S. Army Materiel Command Washington, DC 20315	
13. ABSTRACT			
<p>The closed form inverse of a fairly general tridiagonal matrix is given. The restriction is that the off-diagonal elements in the tridiagonal band be nonzero. If the elements of the matrix are integers, where the upper off-diagonal elements are equal and the lower off-diagonal elements are equal, then an integer multiple of each element of the inverse can be generated by integer arithmetic.</p>			

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

Unclassified

Security Classification

Unclassified

Security Classification

14	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Inverse Tridiagonal Matrix Integer Arithmetic Integer Elements						

II

Unclassified

Security Classification

BALLISTIC RESEARCH LABORATORIES

REPORT NO. 1612

September 1972

THE INVERSE OF A TRIDIAGONAL MATRIX

Palmer R. Schlegel

Vulnerability Laboratory

William Clare Taylor

Applied Mathematics Division

Approved for public release; distribution unlimited.

RDT&E Project No. 1T062110A027

ABERDEEN PROVING GROUND, MARYLAND

BALLISTIC RESEARCH LABORATORIES

REPORT NO. 1612

PRSchlegel/WCTaylor/ats
Aberdeen Proving Ground, Md.
September 1972

THE INVERSE OF A TRIDIAGONAL MATRIX

ABSTRACT

The closed form inverse of a fairly general tridiagonal matrix is given. The restriction is that the off-diagonal elements in the tri-diagonal band be nonzero. If the elements of the matrix are integers, where the upper off-diagonal elements are equal and the lower off-diagonal elements are equal, then an integer multiple of each element of the inverse can be generated by integer arithmetic.

TABLE OF CONTENTS

	Page
ABSTRACT.	3
I. INTRODUCTION.	7
II. FORMS OF MATRIX AND ITS INVERSE	7
III. INTEGER ELEMENTS.	9
REFERENCE	10
DISTRIBUTION LIST	11

I. INTRODUCTION

The closed form inverse of a fairly general tridiagonal matrix is given. The restriction is that the off-diagonal elements in the tridiagonal band be nonzero. If the elements of the matrix are integers, where the upper off-diagonal elements are equal and the lower off-diagonal elements are equal, then an integer multiple of each element of the inverse can be generated by integer arithmetic.

In an earlier note^{1*} an identity was obtained. Here we present a simpler proof of a more general relation. Multiplication of a vector by a tridiagonal matrix consists of applying a three term recurrence operation to the sequence of elements of the vector. We give the explicit expression for the inverse of the matrix in terms of two vectors, U and V, formed by recursive solution of the homogeneous relation with the zero boundary condition at the beginning and at the end respectively.

II. FORMS OF MATRIX AND ITS INVERSE

Let $B = (b_{ij})$ be a tridiagonal matrix such that $b_{i,i+1} \neq 0$ and $b_{i+1,i} \neq 0$, $i = 1, \dots, n-1$. Define $u_0 = 0$, $u_1 = 1$ and $u_{i+1} = -b_{i,i+1}^{-1}(b_{i,i-1}u_{i-1} + b_{ii}u_i)$ for $1 \leq i \leq n-1$ and $v_{n+1} = 0$, $v_n = 1$ and $v_{j-1} = -b_{j,j-1}^{-1}(b_{jj}v_j + b_{j,j+1}v_{j+1})$ for $n+1 > j > 1$. Let

$$\lambda_j^{-1} = b_{j,j-1}v_ju_{j-1} + b_{jj}v_ju_j + b_{j,j+1}v_{j+1}u_j,$$

where

$$\begin{aligned} c_{ij} &= \lambda_j^{-1}v_ju_i, \quad i \leq j \\ &= \lambda_j^{-1}v_iu_j, \quad i > j, \end{aligned}$$

*Superscript numerals refer to references found on page 10.

is the inverse of B . Conversely, if B has an inverse, it is obtained by this construction. To see this let $U^T = (u_1, \dots, u_n)$ and $V^T = (v_1, \dots, v_n)$, where u_i and v_i are defined above. Then $(BU)^T = (0, \dots, 0, \xi)$ and $(BV)^T = (\eta, 0, \dots, 0)$, where ξ and η are some numbers. Let C_j be the j th column of B^{-1} . Since the first $j-1$ elements of BC_j are zero, for some r_j

$$c_{ij} = r_j u_i, \quad i \leq j.$$

Similarly, since the last $n-j-1$ elements of BC_j are zero, for some s_j

$$c_{ij} = s_j v_i, \quad i \geq j.$$

Thus,

$$r_j u_j = c_{jj} = s_j v_j,$$

or for some λ_j

$$r_j = \lambda_j v_j$$

and

$$s_j = \lambda_j u_j.$$

The j th element of BC_j is

$$\begin{aligned} 1 &= b_{j,j-1} c_{j-1,j} + b_{jj} c_{jj} + b_{j,j+1} c_{j+1,j} \\ &= b_{j,j-1} r_j u_{j-1} + b_{jj} r_j u_j + b_{j,j+1} s_j v_{j+1} \\ &= \lambda_j (b_{j,j-1} v_j u_{j-1} + b_{jj} v_j u_j + b_{j,j+1} u_j v_{j+1}), \end{aligned}$$

and this determines λ_j .

III. INTEGER ELEMENTS

Suppose the elements of B are integers. Furthermore, suppose

$b_{i,i+1} = c$ and $b_{i+1,i} = d$ for $i = 1, \dots, n-1$. Let

$$x_{i+1} = -(cdx_{i-1} + b_{ii}x_i),$$

$$y_{j-i} = -(b_{jj}y_j + cdy_{j+1})$$

and

$$w_j = cdx_{j-1}y_j + b_{jj}x_jy_j + cdx_jy_{j+1},$$

where $x_0 = 0$, $x_1 = 1$, $y_{n+1} = 0$ and $y_n = 1$. It can be shown that

$$\begin{aligned} w_j c_{ij} &= c^{j-i} x_i y_j, \quad i \leq j \\ &= d^{i-j} x_j y_i, \quad i > j, \end{aligned}$$

that is, an integer multiple of each element of the inverse can be generated by integer arithmetic.

REFERENCES

1. P. Schlegel, "The Explicit Inverse of a Tridiagonal Matrix," Math of Comp., Vol 24, 1970, p665.